

EXTRACTIVE BIOCONVERSION OF GAMMA-CYCLODEXTRIN GLYCOSYLTRANSFERASE IN ALCOHOL/SALT AQUEOS TWO-PHASE SYSTEM

Tau Chuan Ling, Institute of Biological Sciences, Faculty of Science, University of Malaya,
50603 Kuala Lumpur, Malaysia

Email: tcling@um.edu.my or tauchuan.ling@gmail.com

Yu Kiat Lin, Institute of Biological Sciences, Faculty of Science, University of Malaya,
50603 Kuala Lumpur, Malaysia

Mohamad Suffian Mohamad Annuar, Institute of Biological Sciences,
Faculty of Science, University of Malaya,
50603 Kuala Lumpur, Malaysia

Key Words: aqueous two-phase system, extractive bioconversion, cyclodextrin, *Bacillus cereus*, cyclodextrin glycosyltransferase.

In this study, extractive bioconversion of gamma-cyclodextrin (γ -CD) from soluble starch with *Bacillus cereus* cyclodextrin glycosyltransferase (CGTase, EC 2.4.1.19) was performed in alcohol/potassium phosphate aqueous two-phase system (ATPS). The influences of alcohol-based top phase (ethanol, 1-propanol and 2-propanol) on CGTase bioactivity were investigated. The result showed that the optimum condition was achieved in ATPS composed of 24.0% (w/w) ethanol and 20.0% (w/w) potassium phosphate with 3.0% (w/w) sodium chloride. It demonstrated that the γ -CD was primarily partitioned to the top phase and the CGTase remained in salt-rich bottom phase of the ATPS. The result indicated that a relatively high concentration of γ -CD (2.28 mg/mL) was recovered in the top phase of ethanol/potassium phosphate ATPS after 1 hour bioconversion operation. Continuous repetitive batch (three times) bioconversion of soluble starch was performed successfully.

Extractive bioconversion of gamma-cyclodextrin glycosyltransferase in alcohol/salt aqueous two-phase system

In this study, extractive bioconversion of gamma-cyclodextrin (γ -CD) from soluble starch with *Bacillus cereus* cyclodextrin glycosyltransferase (CGTase, EC 2.4.1.19) was performed in alcohol / potassium phosphate aqueous two-phase system (ATPS). The influences of alcohol-based top phase (ethanol, 1-propanol and 2-propanol) on CGTase bioactivity were investigated. The result showed that the optimum condition was achieved in ATPS composed of 24.0% (w/w) ethanol and 20.0% (w/w) potassium phosphate with 3.0% (w/w) sodium chloride. It demonstrated that the γ -CD was primarily partitioned on the top phase and the CGTase remained at (mg/mL) was recovered in the top phase of ethanol / potassium phosphate ATPS after 1 hour bioconversion operation. Continuous repetitive batch (three times) bioconversion of soluble starch was performed successive.

Key words: aqueous two-phase system, extractive bioconversion, cyclodextrin, *Bacillus cereus*, cyclodextrin, glycosyltransferase.

